

INTERVENTIONAL RADIOLOGY (IR)

MINIMALLY INVASIVE SPECIALISTS

PATIENT INFORMATION

WHAT IS INTERVENTIONAL RADIOLOGY?

Interventional radiology, or IR, is a new subspecialty that has been responsible for major recent medical advancement. It is at the forefront of medical technology and innovation. It is responsible for image-guided minimally invasive procedures and therapies you may have thought carried out by surgeons, but are in fact performed by Interventional radiologists.

IR is a minimally invasive alternative to open surgery that uses radiological image guidance (X-rays, ultrasound, CT and MRI) to aid treatment. For many patients, surgery can be avoided and the risks to the patient can be lowered with faster recovery times. IR can also be used as an adjuvant to surgery and combined procedures with surgeons can optimise patient care.

WHEN DID IR START?

IR developed in diagnostic angiography in the 1960s, after the Dr Sven-Ivar Seldinger described the Seldinger technique to access an artery. The technique used a hollow needle and guide wire to access the artery.

An American radiologist, Dr Charles Dotter, altered the course of cardiovascular interventional radiology in 1964 and is considered the father of IR. He modified the Seldinger technique for therapeutic purposes. IR was born when Dotter dilated a tight stenosis (narrowing) of a superficial femoral artery, using a plastic tube (catheter), in an 82-year-old lady with a painful leg and gangrene due to poor blood supply, who refused amputation. This was all performed through a small 5mm incision in the groin. Following this, despite scepticism from his surgical colleagues, the leg pain ceased, she started walking and her leg improved. Dr Charles Dotter was nominated for the Nobel Prize in Medicine in 1978.

The term 'interventional radiology', however, was not conceived until March 1967 when an American radiologist, Alexander Margulis, described it in an edition of the American Journal of Roentgenology.

WHAT SKILLS DO INTERVENTIONAL RADIOLOGISTS (IRS) HAVE?

IRs have expertise in guiding small needles, catheters and other medical equipment into the body through tiny (5–10 mm) incisions in the skin to treat disease. The treatments performed are truly minimally invasive. However, the basic skills of an IR are still image interpretation and, therefore, core diagnostic radiology is at the heart of interventional radiology training. This combined skill has meant that there is hardly any area of hospital medicine where IR has not had an impact on the patient pathway.

WHAT CAN IR TREAT?

You maybe surprised to know the extent of therapies now performed predominantly by IRs. Traditionally surgeons performed a lot of these procedures, although a significant range of treatments is new and innovative.

Virtually all body parts and systems can be treated using IR techniques. However, broad categories of treatments available include:

1. Draining of fluid collections (abscesses, dilated kidneys, pleural effusions)
2. Opening up of blocked tubes (arteries, veins, bile ducts, ureters, fallopian tubes)
3. Stopping bleeding from any cause (gastrointestinal bleeding, obstetric and gynaecological bleeding, traumatic bleeding, bleeding after surgery, brain aneurysms, vascular malformations) by occluding the vessels with embolisation
4. Destroying tumours in the liver, lung, kidney by delivering local chemotherapy, embolisation or ablation techniques (such as cryotherapy and radiofrequency ablation)
5. Minimally invasive treatments of thoracic and abdominal aneurysms
6. Preoperative embolisation to improve the safety of surgery.

PATIENT SAFETY

IRs pioneered the safe and high-quality procedures and standards for performing image guided minimally invasive therapies.

IRs are specialists of radiology, who have

completed further education and training in diagnostic radiology and interventional radiology including radiation safety, radiation physics, the biological effects of radiation, injury prevention and clinical practice.

WHAT ARE THE ADVANTAGES OF IR?

IR therapies are minimally invasive and typically only require a short stay in hospital. A significant number of these procedures are performed as day cases. General anaesthesia is usually not required. Risk, pain and recovery time are usually reduced compared with conventional surgery.

CAN GPs REFER DIRECTLY TO AN IR?

Interventional radiology is a growing subspecialty in which we cater for all aspects of the patient's pathway. We perform outpatient clinics, pre-admission assessments and have access to day-case beds if you need a simple procedure e.g. varicocele embolisation, uterine artery embolisation or paracentesis.

The IR's at Queen Elizabeth Hospital Birmingham offer direct referrals for image guided minimally invasive procedures from GPs. There is no need to be seen by another specialist within the hospital. If procedures were more complex and need in-patient care then we often would admit patients as part of a multidisciplinary team.

IR's typically work as part of a multidisciplinary team within hospitals, so are always able to refer you to another team member if it is appropriate.

IR's are happy to discuss your patient's requirements and give advice regarding the most appropriate radiological investigations and treatments available.

MILESTONES OF IR

You may be surprised to learn of the advancements in medicine that have been primarily down to IR innovation. You may not even have known that IR had been involved.

Here are a few milestones of IR

- 1964 Angioplasty
- 1966 Embolisation of spinal tumour
- 1967 Judkins technique for coronary angiography
- 1969 Prototype catheter delivered stent
- 1970s Embolisation coils
- 1972 Embolisation of bleeding gut
- 1973 Embolisation of pelvic bleeding secondary to trauma
- 1974 Selective arterial thrombolysis
- 1977 Embolisation of pulmonary arteriovenous malformations
- 1978 Embolisation of varicoceles
- 1980s Biliary stents
- 1982 Transjugular intrahepatic portosystemic shunts (TIPS)
- 1983 development of balloon expandable stents
- 1990s Treatment of bone and kidney tumours by embolisation
- 1990 Radiofrequency ablation of liver tumours
- 1991 Endovascular stent grafts to treat aortic aneurysms

- 1995 Uterine artery embolisation to treat fibroids (UAE)
- 1999 Endovascular laser therapy (EVLV) to treat varicose veins.

CONTACT INFORMATION:

THE RADIOLOGY CLINIC

TEL: 0121 455 9496

EMAIL

info@ianmccafferty.com



BRITISH SOCIETY OF INTERVENTIONAL RADIOLOGY

www.bsir.org

This leaflet has been prepared by the British Society of Interventional Radiology (BSIR) and the Clinical Radiology Patients' Liaison Group (CRPLG) of the Royal College of Radiologists. Approved by the board of the Faculty of Clinical Radiology: 25 February 2011.

Legal notice

Please remember that this leaflet is intended as general information only. It is not definitive, and the RCR and the BSIR cannot accept any legal liability arising from its use. We aim to make the information as up to date and accurate as possible, but please be warned that it is always subject to change. Please therefore always check specific advice on the procedure or any concerns you may have with your doctor.